

Arid Zone Trees



Dodonaea angustifolia *Hopbush*



Horticultural Qualities **Dodonaea angustifolia** *Hopbush*

Foliage: Evergreen
Mature Height: 12' - 20'
Mature Width: 10' - 15'
Growth Rate: Moderate
Hardiness: 18 degrees F
Exposure: Full Sun
Leaf Color: Green
Shade: Filtered
Flower Color: Yellow-Green
Flower Shape: Petal-less
Flower Season: Spring to Fall
Thorns: None
Box Sizes Produced: 24"
Propagation Method: Seed



www.aridzonetrees.com

Arid Zone Trees, P. O. Box 167, Queen Creek, AZ 85242, Phone 480-987-9094 e-mail: info@aridzonetrees.com

Dodonaea angustifolia

Hopbush

Dodonaea angustifolia (Hopbush) is a evergreen shrub form measuring in height from 12 to 20 feet and width from 10 to 15 feet. Its native habitat is on rocky slopes in the lower elevations of the Sonoran Desert to the Oak woodland in the higher elevations of 6000 ft. It is frost sensitive (little leaf damage) at 25 degree F, however survives at 18 degrees F. Inconspicuous flowers are wind pollinated and the winged fruit is also dispersed in the wind.



Cultural Practices

Foster the development of a more dispersed root system and reduce the risk of wind throw by arranging irrigation emitters at varying distances from the trunk to encourage roots to "seek out" water and nutrients. Irrigation emitter arrangement along with other information on irrigations practices for desert trees can be found at www.aridzonetrees.com and click on the FAQ link.

Prune as needed to reinforce the structure and form of the tree. Periodic thinning is the most desirable method of pruning. Avoid hedging or heading back desert species, as this will only stimulate excessive branching. Do not remove more than 30% of the canopy during the summer as this can lead to sunburn injuries that can later be invaded by wood boring insects. Always use clean, sharp tools that are cleaned regularly in a 10% solution of bleach. For detail pruning guide see www.aridzonetrees.com and click on the FAQ interactive button.

Periodically insect pests can be a problem on some desert trees. On young trees, insect infestation can slow typical seasonal growth. Inspect trees during the growing season for common garden sucking insects such as aphids, thrip, whiteflies or psyllids. During dry months, (May and June) in dusty conditions, spider mites can appear. Monitor for infestation and apply controls as needed. Spray applications of water or water and Safer Soap give short-term control (3 to 7 days) for small insect population. For heavy infestation or longer control use federally registered insecticides. A contact insecticide application will kill existing adults. An application with a systemic soil drench will provide 8 to 12 weeks control for any post application insect hatchings or migration of insects. Before using pesticide for the first time or on new plants or cultivar, treat a few plants and check for phytotoxicity. **Always read label and follow label instruction before using pesticides. For pesticide control recommendations contact a licensed pest control advisor.**

www.aridzonetrees.com

Arid Zone Trees, P. O. Box 167, Queen Creek, AZ 85242, Phone 480-987-9094 e-mail: info@aridzonetrees.com